



Contents lists available at ScienceDirect

Journal of Acute Disease

journal homepage: [www.jadweb.org](http://www.jadweb.org)

Original article doi:

©2016 by the Journal of Acute Disease. All rights reserved.

## Characterization of integrons and associated gene cassettes in *Acinetobacter baumannii* strains isolated from intensive care unit in Tehran, Iran

Hossein Goudarzi<sup>1</sup>, Mehdi Azad<sup>2</sup>, Sima Sadat Seyedjavadi<sup>3</sup>, Hadi Azimi<sup>4</sup>, Alireza Salimi Chirani<sup>1</sup>, Vahid Fallah Omrani<sup>5</sup>, Mehdi Goudarzi<sup>1\*</sup><sup>1</sup>Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran<sup>2</sup>Department of Medical Laboratory Sciences, School of Paramedicine, Qazvin University of Medical Sciences, Qazvin, Iran<sup>3</sup>Department of Medical Mycology, Pasteur Institute of Iran, Tehran, Iran<sup>4</sup>Department of English Language Teaching, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran<sup>5</sup>Cellular and Molecular Biology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

### ARTICLE INFO

#### Article history:

Received 23 Jun 2016

Received in revised form 30 Jun 2016

#### Keywords:

*Acinetobacter baumannii*

Multidrug resistance

Integron

Intensive care unit

### ABSTRACT

**Objective:** To determine the antimicrobial susceptibility patterns, the frequency of integrons and associated gene cassettes in *Acinetobacter baumannii* (*A. baumannii*) strains isolated from selected hospital intensive care units.

**Methods:** During a ten-month period, 120 *A. baumannii* isolates were studied. The resistance rates to different classes of antimicrobial agents were determined. PCR was used to detect different types of integrons and associated gene cassettes.

**Results:** The resistance rates to the majority of antibiotics tested were found to be between 39.3% and 99.1%. No isolate was observed to be resistant to colistin and polymyxin B. The rate of extensive drug-resistance among these clinical isolates was 62.5%. The prevalence of class 1 and 2 integrons was found to be 74.1% and 12.5%, respectively. Seven different gene cassettes (*ampC*, *aacA4-catB8*, *ISAbal1-bla<sub>OXA-23</sub>-GES-14*, *aadA2-cm1A6-GES-14-qacF*, *VIM-25-GES-24-qacF*, *dfrA5-ISAbal1-bla<sub>OXA-51</sub>-bla<sub>OXA-40</sub>* and *aadA2-GES-11-IMP-1*) were observed in Class 1 integron-carrying strains. Three gene cassettes (*IMP-4*, *VIM-2-VEB-aacA4* and *dfrA2-sat-2-aadA4*) were detected in class 2 integron-bearing *A. baumannii* strains.

**Conclusions:** A high prevalence of integron was described among multidrug resistant *A. baumannii* in the hospital. The findings highlighted the need for continuous surveillance in order to prevent dissemination of multidrug resistance among *A. baumannii* strains in Iran.

## 1. Introduction

*Acinetobacter baumannii* (*A. baumannii*), as an important nosocomial pathogen especially in intensive care units (ICUs), is responsible for a wide range of infections that can be ranged from urinary tract infections to surgical wounds infection, ventilator-associated pneumonia, meningitis, bacteremia, and life threatening infections. The most important factor contributing to the successful

extensive distribution of this nosocomial pathogen is stated to be its remarkable ability for the acquisition of a wide variety of antibiotic resistance genes and also adaptation in various harsh environments[1]. The acquisition of a wide variety of antibiotic resistance genes not only leads to an increase in economic burden, but also causes serious therapeutic problems. Moreover, it can lead to difficulties in infection control in hospitals and eradication of the bacteria. The emergence and extremely rapid spread of multidrug resistant *A. baumannii* isolates are becoming a serious concern in global public health. The spread of the resistant genes in hospitals and community is mediated by horizontal gene transfer[2]. Mobile elements including plasmids, transposons, or integrons are the most efficient genetic elements promoting acquisition and dissemination of resistance determinants[3]. During the past several decades, despite introduction of new therapeutic options, *A. baumannii* strains have shown a remarkable ability to rapidly develop multidrug resistance (MDR). This rapid increase of MDR is not only due to

\*Corresponding author: Mehdi Goudarzi, Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Tel/Fax: +98 (21) 23872556

E-mail: [gudarzim@yahoo.com](mailto:gudarzim@yahoo.com)

The study protocol was performed according to the Helsinki declaration and approved by Ethics Committee of Shahid Beheshti University of Medical Sciences, Tehran, Iran (No. 13478).

Foundation Project: Supported by the grant from research deputy of Shahid Beheshti University of Medical Sciences (grant number?).

The journal implements double-blind peer review practiced by specially invited international editorial board members.